

^a Key Laboratory of Laboratory Medicine of Jiangsu Province, Medical School, Jiangsu University, Zhenjiang, Jiangsu, P. R. China; ^b The Affiliated Hospital, Jiangsu University, Zhenjiang, Jiangsu, P. R. China

Contact: Wenrong Xu, Ph D. M D, Professor, Medical School, Jiangsu University, 301 Xuefu Road, 212013, Tel: +86 511 85038215, Fax: +86 511 85038483, E-mail: ics@ujs.edu.cn; Hui Qian, Ph D. M D, Professor, Medical School, Jiangsu University, 301 Xuefu Road, 212013, Tel: +86 511 85038334, Fax: +86 511 85038483, E-mail: lstmmlst@163.com; ^c These authors contributed equally to this work

Received March 10, 2014; accepted for publication June 11, 2014; available online without subscription through the open access option.

©AlphaMed Press
1066-5099/2014/\$30.00/0

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/stem.1771

HucMSC-exosome Mediated -Wnt4 Signaling is Required for Cutaneous Wound Healing

BIN ZHANG ^{a,c}, MEI WANG ^{a,c}, AIHUA GONG ^{a,c}, XU ZHANG ^a, XIAODAN WU ^a, YANHUA ZHU ^a, HUI SHI ^a, LIJUN WU ^a, WEI ZHU ^a, HUI QIAN ^{a,*}, WENRONG XU ^{a,b*}

Key words. Exosomes • mesenchymal stem cells • wound healing • Wnt4 • β -catenin • AKT.

ABSTRACT

Mesenchymal stem cell-derived exosomes (MSC-Ex) play important roles in tissue injury repair, however, the roles of MSC-Ex in skin damage repair and its mechanisms are largely unknown. Herein, we examined the benefit of human umbilical cord mesenchymal stem cell derived exosome (hucMSC-Ex) in cutaneous wound healing using a rat skin burn model. We found that hucMSC-Ex-treated wounds exhibited significantly accelerated re-epithelialization, with increased expression of CK19, PCNA, collagen I (compared to collagen III) in vivo. HucMSC-Ex promoted proliferation and inhibited apoptosis of skin cells after heat-stress in vitro. We also discovered that Wnt4 was contained in hucMSC-Ex, and hucMSC-Ex derived Wnt4 promoted β -catenin nuclear translocation and activity to enhance proliferation and migration of skin cells, which could be reversed by β -catenin inhibitor ICG001. In vivo studies confirmed that the activation of Wnt/ β -catenin by hucMSC-Ex played a key role in wound re-epithelialization and cell proliferation. Furthermore, knockdown of Wnt4 in hucMSC-Ex abrogated β -catenin activation and skin cell proliferation and migration in vitro. The in vivo therapeutic effects were also inhibited when the expression of Wnt4 in hucMSC-Ex was interfered. In addition, the activation of AKT pathway by hucMSC-Ex was associated with the reduction of heat stress-induced apoptosis in rat skin burn model. Collectively, our findings indicate that exosome-delivered Wnt4 provides new aspects for the therapeutic strategy of MSCs in cutaneous wound healing. *STEM CELLS 2014; 00:000–000*

INTRODUCTION

Cutaneous wound requires a well-orchestrated integration of the differentiation, migration, proliferation and apoptosis of skin cells to create the multilayered tissue that constitutes the skin [1-3]. Skin burn is very common and has a consistently high rate of mortality and morbidity [4], which not only destroy the barrier function of the skin but also alter the perceptions of pain, temperature and touch [5]. Thus, there is an urgent

need to find an alternative approach to prompt wound healing.

MSCs have a significant promise for regenerative medicine due to convenient isolation, low immunogenicity, the ability to transdifferentiate and create a favorable environment for tissue regeneration [6, 7]. HucMSCs are inexhaustible and can be harvested at a low cost without an invasive procedure and they are promising cells in the formation of tissue regenerative during inflammation and tissue injuries [8, 9]. We have

previously found that hucMSCs can ameliorate acute kidney injury (AKI) and mouse hepatic injury [10-13]. Interestingly, Bruno and colleagues confirmed that microvesicles derived from MSCs could protect against acute renal tubular injury [14, 15]. Similarly, MSCs derived exosomes (MSC-Ex) have been proved to contribute to neurite outgrowth [16], which has been described as a new mechanism of cell-to-cell communication [17]. Recently, we also found that hucMSC-Ex alleviated liver fibrosis and promoted renal injury repair [18, 19]. Therefore, it is suggested that MSC-Ex might be a main mechanism of MSCs effects on wound healing. Since exosomes delivered components are complicated, exosomes-mediated signaling might have more differences in various injury models. Wnt/ β -catenin signaling plays an important role in skin development [3, 20-22] and wound healing [23]. The active Wnt proteins can be delivered by exosomes and affect gene expression of target cells [24, 25]. However, whether hucMSC-Ex mediated delivery of Wnts and its function in wound healing are not clear.

In this study, we investigated the role of hucMSC-Ex in wound healing by using deep second-degree burn injury. HucMSC-Ex-derived Wnt4 protein activates β -catenin signaling in skin cells and promotes their proliferation and migration, leading to the enhancement of wound healing. Moreover, our findings indicate that heat stress inhibits AKT signaling in skin cells, while treatment with hucMSC-Ex reverses this inhibition and improves the survival of skin cells.

MATERIALS AND METHODS

The study was approved by the ethical committee of Jiangsu University (2012258).

Cell culture

HucMSCs were isolated and identified as previously described [26]. HucMSCs and human lung fibroblasts (HFL1) were cultured in serum-free DMEM medium (Life Technologies). Keratinocytes HaCAT cells were purchased from ATCC and maintained in DMEM containing 10% fetal bovine serum (FBS; Gibco, Grand Island, USA) at 37°C with 5% CO₂. Dermal fibroblasts (DFL) were isolated and sorted from rat back skin as previously described [27, 28] and cultured in DMEM containing 10% FBS at 37°C with 5% CO₂.

Isolation and characterization of exosomes

Exosomes were extracted and purified as previously described [18, 19]. Cell supernatants were centrifuged to remove cell debris and then passed through a 0.22- μ m filter. Final exosomes were obtained and stored at -70°C. The protein content, as the quantification of exosomes, was determined by using a BCA protein assay kit (CW BIO, Shanghai, China). The final concentration of exosomes used for treating skin cells in vitro was 160 μ g/ml and total 200 μ g exosomes were applied to

treat each animal. The morphology of the extracted exosomes was observed by using transmission electron microscopy (FEI Tecnai 12, Philips, Netherlands). Size distribution within exosome preparations was analyzed by measuring the rate of Brownian motion using a NanoSight LM10 system which is equipped with a fast video capture and particle-tracking software (NanoSight, Amesbury, UK). The CD9, CD63, and CD81 molecules which frequently located on the surface of exosomes were analyzed by using Western blot.

Rat skin wound model and treatment

Adult female Sprague-Dawley rats (weighing 220 \pm 20 g) were purchased from the Animal Centre of Chinese Academy of Sciences (Shanghai, China). Rat model of skin deep second-degree burn wound was established as described previously with added modifications [29]. Rats were anesthetized with sodium thiopental at a dose of 40 mg/kg body weight. After the hair on their upper back was shaved, the back skin of rats were injured with 80 °C water for 8 seconds to create a 16 mm diameter wound, then covered with gauze soaking saline for 6 minutes on the wound. Meanwhile, 1 \times 10⁶ cells (hucMSC and HFL1) suspended in 200 μ l PBS, 200 μ g exosome (hucMSC-Ex and HFL1-Ex) suspended in 200 μ l PBS, or 200 μ l PBS were injected subcutaneously at three sites. The normal group had no treatment. The animals were housed individually. At 1 week and 2 weeks after treatment, the rats were sacrificed and the wound area was cut for further analysis.

H&E staining

The wound skin and surrounding skin (4 mm²) were fixed in 4% paraformaldehyde (pH 7.4), and gradually dehydrated, embedded in paraffin, cut into 4- μ m sections and stained with H&E stain for light microscopy. Each slide was given a histological score ranging from 1 to 10 according to the following parameters [30]. The criteria used for histological scores of wound healing are summarized in **Supplementary table 1**.

Wnt reporter activity assay

For the luciferase reporter assay, HEK293T cells were co-transfected with TOP-Flash or FOP-Flash luciferase reporter. Transfection efficiency was normalized by co-transfection with a β -actin-Renilla reporter containing a Renilla luciferase gene under the control of a human β -actin promoter. The activities of firefly luciferase and Renilla luciferase were quantified by using the dual-luciferase reporter assay system (Promega). For quantification of Wnt activity, after transfection for 6 hours, HEK293T cells were treated with equal amounts of exosomes or PBS from different samples. Wnt reporter activity was determined by TOP / FOP luciferase.

Lentiviral knockdown of Wnt4 in hucMSCs

The lentiviral expression vector containing the Wnt4 shRNA sequence (sigma) was selected for specifically

targeting Wnt4 silence, which was classified as Lenti-wnt4-shRNA, and Lenti-GFP-shRNA as negative control vector. The Wnt4 shRNA lentivirus vectors were generated by ligating the vector Tet-pLKO-puro. Wnt4 shRNA oligonucleotide sequences are: Forward, 5'-CCGGCCCAAGAGATACTGGTTGTATCTCGAGATACAACCA GTATCTCTGGGTTTTG-3'; Reverse, 5'-AATTCAAAAACCAAGAGATACTGGTTGTATCTCGAGATAC AACCAGTATCTCTGGG-3'. The sequences of control shRNA are: Forward, 5'-CCGGGCAAGCTGACCCTGAAGTTCATCTCGAGATGAACCTC AGGGTCACGTTGCTTTTTG-3' Reverse, 5'-AATTCAAAAAGCAAGCTGACCCTGAAGTTCATCTCGAGAT GAACCTCAGGGTCACGTTG-3'. The recombinant lentivirus was produced by co-transfecting HEK293T cells with pLKO-GFP-shRNA or pLKO-Wnt4-shRNA, PU1562 and PU1563 plasmid by using Lipofectamine 2000 (Invitrogen). The virus-containing supernatant was harvested at 48 h and 72 h post-transfection. HucMSCs were transduced with the prepared lentivirus (Lenti-wnt4-shRNA or Lenti-GFP-shRNA). Stable cell lines were obtained after selection with 1 µg/mL of puromycin (Invitrogen) for 15 days. The expression of shRNA was induced by addition of 80 µg/mL doxycycline for 2 days. The efficiency of wnt4 knockdown was evaluated by using real-time quantitative RT-PCR and Western blot. The stable cell lines were cultured in serum-free medium for 48 h, then the supernatants were collected and exosomes isolated for further study.

TUNEL assay

The apoptotic skin cells in tissue slides were measured by using an in situ cell apoptosis detection kit according to the manufacturer's instruction (Boster). The number of positive cells was calculated in three or more random fields.

Statistical analysis

All data were shown as means ± standard deviation (SD). The statistically significant differences between groups were assessed by analysis of variance (ANOVA) or t-test using Prism software (GraphPad, San Diego, USA). P value <0.05 was considered significant.

RESULTS

HucMSC-Ex promotes cell proliferation and re-epithelialization in rat deep second-degree burn injury model

To investigate the roles of hucMSC-Ex in wound healing, hucMSC-Ex was firstly extracted and identified as previously described [18, 19]. The morphology of hucMSC-Ex was observed under transmission electron microscopy, and its size was measured by using NanoSight analysis. The results showed that hucMSC-Ex was about 100 nm spherical vesicles (Fig. S1A and S1B). The results of Western blot showed that exosomal markers, including CD63, CD81, and CD98 were expressed in both hucMSC-

Ex and control exosomes from human lung fibroblasts HFL1 (HFL1-Ex) (Fig. S1C).

We established a rat deep second-degree burn injury model and infused exosomes and their derived cells into the injured rats, separately. The results of histological evaluation of wounds at 1 week post-infusion showed that the number of epidermal and dermal cells significantly increased in hucMSCs or hucMSC-Ex treated wounds (Fig. 1A), while wounds that were treated with PBS, HFL1 cells or HFL1-Ex were still in second-degree burn injury state (Fig. 1A). The results of PCNA immunochemical staining showed that hucMSCs and hucMSC-Ex groups had more PCNA-positive cells in wound area than that in PBS, HFL1, and HFL1-Ex groups at both 1 and 2 weeks post-infusion (Fig. 1B).

Strikingly, at 2 weeks post-infusion, HE staining results showed that wounds treated with hucMSCs or hucMSC-Ex significantly enhanced re-epithelialization (complete re-epithelialization in all 6 wounds; n=6) compared to the PBS group (complete re-epithelialization in 1 of 6 wounds; n=6) or HFL1 group (complete re-epithelialization in 2 of 6 wounds; n=6) or HFL1-Ex (complete re-epithelialization in 2 of 6 wounds; n=6) (Fig. 1A). In consistent with the above results, the histological scores were significantly higher in hucMSC and hucMSC-Ex groups than that in PBS, HFL1 and HFL1-Ex groups (Fig. 1C). We determined the relative expression of collagen I to collagen III by using quantitative RT-PCR and found that the ratio of collagen I to collagen III was higher in hucMSC and hucMSC-Ex groups than that in PBS, HFL1 and HFL1-Ex groups, suggesting that hucMSCs and hucMSC-Ex are able to reduce the formation of scar in wound area (Fig. 1D).

To confirm the promoting role of hucMSC-Ex in re-epithelialization, we determined the expression of CK19, a epithelial marker, by using immunofluorescent staining. The results revealed that CK19 expression was remarkably higher in hucMSC and hucMSC-Ex groups at 1 week post-infusion than that in PBS, HFL1 and HFL1-Ex groups (Fig. 1E). At 2 weeks post-infusion, there formed a complete epidermal structure in the CK19-positive area of the wounds in hucMSC and hucMSC-Ex groups but not in PBS, HFL1 and HFL1-Ex groups (Fig. 1E). The increased levels of CK19 and PCNA in hucMSC and hucMSC-Ex groups were further confirmed by using Western blot (Fig. 1F). Taken together, these results indicate that hucMSC-Ex prompt the same repair of skin second-degree burn injury as hucMSCs by enhancing proliferation of skin cells and re-epithelialization in wound area.

HucMSC-Ex inhibits heat stress-induced apoptosis and enhances proliferation of skin cells in vitro

In order to verify the above results and explore the mechanisms for hucMSC-Ex-induced repair, we treated the immortal human keratinocytes HaCAT and primary cultured dermal fibroblasts (DFL) at 43°C for 40 min to

mimic burn injury model *in vivo*. In comparison to PBS group, hucMSC-Ex significantly inhibited heat stress-induced apoptosis in HaCAT while HFL1-Ex had minimal effect. The similar effects were also observed in primary DFL (Fig. 2A). The results of cell proliferation assay showed that treatment with hucMSC-Ex but not HFL1-Ex promoted the proliferation of HaCAT and DFL cells in a time-dependent manner after heat stress (Fig. 2B and 2C). The expression of PCNA was enhanced by hucMSC-Ex but not HFL1-Ex in HaCAT and DFL cells (Fig. 2D and 2E). Western blot analyses of apoptosis-associated proteins in HaCAT and DFL cells showed that Bax level was lower in hucMSC-Ex group than that in PBS and HFL1-Ex groups, while the expression of Bcl-2 was higher in hucMSC-ex group than that in PBS and HFL1-Ex groups (Fig. 2D and 2E). These results suggest that hucMSC-Ex inhibits heat stress-induced apoptosis in skin cells and prompts their proliferation.

HucMSC-Ex activates Wnt/ β -catenin signaling to prompt wound healing

Considering the significant role of β -catenin signaling in skin development [3, 20-22] and cutaneous wound healing [23], we hypothesized that wnt/ β -catenin signaling might be involved in the biological effects of hucMSC-Ex on wound healing. We found that hucMSC-Ex enhanced TOP-flash reporter activity in 293T cells, while HFL1-Ex had no effect (Fig. 3A). HucMSC-Ex treatment induced more nuclear translocation of β -catenin in HaCAT and DFL cells than PBS and HFL1-Ex (Fig. 3B and S2A). The expression of β -catenin and its downstream genes (cyclin-D1, cyclin-D3, and N-cadherin) in DFL cells were significantly increased by hucMSC-Ex but not PBS and HFL1-Ex (Fig. S2B). The induction of β -catenin downstream genes (cyclin-D1, cyclin-D3, N-cadherin) and the increase of TOP-flash reporter activity by hucMSC-Ex were almost completely abrogated by ICG001, which selectively inhibits β -catenin/CBP interaction (Fig. 3C and 3D), suggesting the specific activation of β -catenin signaling by hucMSC-Ex.

To demonstrate the functional role of β -catenin activation by hucMSC-Ex, we detected the cell cycle and migration of skin cells. The results showed that hucMSC-Ex significantly promoted wound closure of scratch, which could be reversed by simultaneous treatment with ICG001 (Fig. 3E and S2C). HucMSC-Ex prompted the transition of G1 to M phase during cell cycle (Fig. 3F). We next determined the role of β -catenin activation in hucMSC-Ex-mediated wound healing *in vivo*. Co-injection of ICG001 significantly inhibited the enhancement of wound healing and the increase of PCNA expression by hucMSC-Ex *in vivo* (Fig. 3G and 3H). In summary, hucMSC-Ex activates wnt/ β -catenin signaling to enhance wound healing.

HucMSC-Ex delivered-Wnt4 induces β -catenin activation

Since hucMSC-Ex activated Wnt/ β -catenin signaling to prompt wound healing, we next focused on which component in hucMSC-Ex that mediated this effect. We screened the expression of Wnt family members including Wnt1, Wnt2, Wnt3, Wnt3a, Wnt4, Wnt5a, Wnt6, Wnt7b, Wnt10b and Wnt11 in hucMSCs and HFL1 cells. We found that Wnt4 was significantly higher in hucMSCs than that in HFL1 cells (Fig. 4A). The higher expression of Wnt4 in hucMSC-Ex was further confirmed by using Western blot (Fig. 4B). To investigate the role of Wnt4 in hucMSC-Ex-mediated β -catenin activation and wound healing, we knocked down Wnt4 in hucMSCs by using shRNA (Fig. 4C). Knockdown of Wnt4 reduced the expression of Wnt4 in hucMSC-Ex and inhibited the transcriptional activity of β -catenin in 293T cells (Fig. 4D). The enhanced nuclear translocation of β -catenin and the increased expression of its downstream targets by hucMSC-Ex were also abrogated by Wnt4 knockdown (Fig. 4E and 4F; Fig. S3A and S3B). The phosphorylation of GSK3 β , a classical negative regulator of Wnt signaling pathway, was promoted by hucMSC-Ex. However, knockdown of Wnt4 in hucMSC-Ex barely affected this phenomenon (Fig. 4F).

Furthermore, hucMSC-Ex-induced cell cycle progression and cell migration were also abolished by Wnt4 knockdown (Fig. 4G and 4H; Fig. S3C). Wnt4 knockdown delayed wound healing and reduced the expression of PCNA induced by hucMSC-Ex *in vivo* (Fig. 4I and 4J). These results indicate that Wnt4 plays an important role in hucMSC-Ex-mediated wound healing.

HucMSC-Ex reverses acute thermal injury-induced apoptosis in skin cells through activation of AKT pathway

In order to explore the mechanism by which hucMSC-Ex reversed acute thermal injury-induced apoptosis *in vitro* and *in vivo*, we determined the status of AKT and MAPK signaling in HaCAT and DFL treated with or without hucMSC-Ex after heat stress. There was no significant change in MAPK pathway except a slight down-regulation of phosphorylated p38 after hucMSC-Ex treatment (Fig. S4). However, the phosphorylated form of AKT was significantly increased in both HaCAT and DFL cells after hucMSC-Ex treatment (Fig. 5A). AKT inhibitor LY294002 suppressed the activation of AKT and the induction of Bcl-2 by hucMSC-Ex (Fig. 5B).

In consistent with the *in vitro* study, the activation of AKT and the increase of Bcl-2 protein level by hucMSC-Ex was also observed in skin tissues by using Western blot and immunofluorescent staining (Fig. 5C and 5D). Moreover, hucMSC-Ex treatment reduced the number of apoptotic cells in wound area while the simultaneous treatment with LY294002 reversed this effect (Fig. 5E). We further tested the role of AKT signaling in hucMSC-Ex-mediated promotion of cell proliferation and migration. As shown in Fig. 5F, hucMSC-Ex promot-

ed cell cycle transition from G1 to S phase, which could be partially inhibited by LY294002, suggesting that other pathways may also participate in this process. The similar effect of LY294002 on hucMSC-Ex-induced cell migration was also observed (Fig. 5G). These results suggested that hucMSC-Ex reversed acute thermal injury-induced apoptosis in skin cells mainly through activation of AKT pathway. To explore input signal of AKT pathway via hucMSC-Ex, we analyzed the cytokines within huc-MSC exosomes that may activate AKT pathway by luminex assay. The data indicated that hucMSC-Ex could deliver many cytokines including PDGF-BB, G-CSF, VEGF, MCP-1, IL-6 and IL-8, which may activate AKT signaling (Supplementary table 5).

HucMSC-Ex mediates bifurcated activation of Wnt4/ β -catenin and AKT signaling to promote wound healing

The Wnt/ β -catenin signaling can be directly or indirectly regulated by AKT [31, 32]. Considering that AKT and β -catenin signaling were both activated by hucMSC-Ex, we next investigated the relationship between AKT and β -catenin signaling after hucMSC-Ex treatment. Inhibition of AKT by LY294002 had minimal effects on the increased TOP-flash reporter activity and the nuclear translocation of β -catenin induced by hucMSC-Ex (Fig. 6A and 6B), but reversed the phosphorylation of GSK3 β induced by hucMSC-Ex (Fig. 6C), suggesting that exosomal Wnt4 mediates the activation of Wnt/ β -catenin signaling independent of GSK3 β . In consistent with the *in vitro* results, LY294002 almost completely inhibited the activation of AKT but had minimal effect on the increase of β -catenin by hucMSC-Ex *in vivo* (Fig. 6D). In contrary, ICG001 treatment and knockdown of Wnt4 had no effect on the activation of p-AKT by hucMSC-Ex (Fig. 6E and 6F). The above results revealed that hucMSC-Ex enhanced wound healing by parallel activation of Wnt4/ β -catenin and AKT signaling.

DISCUSSION

In the present study, we investigated whether hucMSC-Ex had the same therapeutic effect as hucMSCs in skin injury models and explored the underlying mechanisms. We demonstrated that hucMSC-Ex significantly promoted wound healing in a rat deep second-degree burn injury rat model.

Exosomes are emerging as a new mechanism for cell-to-cell communication and is an important part of cells [14]. Exosomes are known to contain mRNAs, microRNAs and proteins [33, 34]. The previous study shows MSC-Ex promote tissue injury repair through the horizontal transfer of mRNAs and microRNAs [15, 17, 35]. The active proteins can be delivered by exosomes resulting in biological effects of target cells [36, 37]. But few studies have defined the function of protein molecules transported by MSC-Ex in regenerative medicine.

Accumulating evidence indicates that Wnts require a particular lipid modification for proper secretion and function. Wnts are lipid-modified by the acyltransferase porcupine in the endoplasmic reticulum and acts on target cells in an autocrine or paracrine manner [38, 39]. The unmodified Wnts are tightly associated with the plasma membrane and are hard to spread over a distance to act on target cells [40, 41]. In tissue culture cells that are stably expressing Wingless (Wg; *Drosophila* ortholog of Wnt1), only a fraction of the Wnts is actually secreted into the media [42]. In mammalian and *Drosophila* cells, Wnts can not be detected in exosome-free cell culture media [24]. Interfering with the lipid modification of Wnts results in a protein that is no longer active or hydrophobic [42-44]. In addition to essential lipid modification, proper secretory paths of Wnts are required for their function [45]. Several mechanisms have been proposed to explain how Wnts might function as long-range signaling molecules, including their lateral diffusion by association with heparan sulfate proteoglycans (HSPGs), solubilization by high-density lipoproteins and carrier proteins [40, 41]. Recently, several groups confirm that exosomes are important carriers for Wnt secretion and extracellular traveling [24, 25, 46]. Exosomes-mediated delivery of Wnts promotes the growth of *Drosophila* wings [24]. In this study, we found that hucMSC-Ex can deliver Wnt4 to enhance wound healing. Wnt4 knockdown in hucMSC abrogated β -catenin activation skin cells *in vitro* and the therapeutic effects of hucMSC-Ex *in vivo*, suggesting that hucMSC-exosomal Wnt4 is critical for the biological activities of hucMSC-Ex in wound healing.

Although our results showed that hucMSC exosomes promoted the phosphorylation of GSK3 β and inhibited the activity of GSK3 β . However, knockdown of Wnt4 in hucMSC-Ex barely affected the activity of GSK3 β . Recent study has revealed that exosome-free Wnt signaling sequester GSK3 β inside multivesicular endosomes to indirectly inhibit its activation [47]. It has been shown that AKT activation results in the phosphorylation of GSK3 β [32]. Our data revealed that hucMSC exosome delivered many factors, such as G-CSF, PDGF-BB, VEGF, MCP-1, IL-6, IL-8, can affect the activation of AKT/GSK3 β pathway. We further discovered that PI3K/AKT pathway by hucMSC-Ex also partly affected the activation of Wnt/ β -catenin signaling through inhibiting the activity of GSK3 β , which synergized with exosomal Wnt4. However, the interference experiments of Wnt4 confirmed hucMSC-Ex mediated Wnt4 was the key factor for activating β -catenin signaling. That may be why AKT pathway inhibitor LY294002 could slightly inhibit transcription activity of β -catenin and the promoting role of hucMSC-Ex in skin cells migration and cell cycle transition were not completely reversed by LY294002. Because the exosomal components are very complicated, it requires further studies to determine whether there are other molecules enhancing the activity of Wnt/ β -catenin signaling in wound healing.

Previous studies have revealed that apoptosis plays an important role in regulating stem cell-dependent regeneration at the early stages of wound healing [48, 49]. Thus, it is necessary to investigate the mechanism of anti-apoptotic effects by hucMSC-Ex. Further study showed the knockdown of Wnt4 and ICG001 treatment did not significantly affect the expression of apoptosis-related proteins Bcl-2 and Bax. This result indicated that hucMSC-Ex may inhibit apoptosis through other pathway(s). Long-term heat stress (4 weeks) can promote AKT activation and proliferation of fibroblasts [50]. Deregibus et al. reported that endothelial progenitor cell-derived microvesicles activate AKT signaling by a horizontal transfer of mRNAs [51]. Our results indicated that acute thermal injury inhibited AKT signaling and can be reversed by hucMSC-Ex. The results of Luminex assay showed that hucMSC-Ex contained many soluble factors, which have been reported to affect AKT pathway. However, due to the complexity of hucMSC exosomal components, it is hard to determine which factor(s) in hucMSC-Ex mediate the activation of AKT pathway.

In summary, we find the beneficial role of hucMSC-Ex in wound healing by using rat burn injury. HucMSC-Ex enhances wound closure by delivering Wnt4 to activate Wnt/ β -catenin in skin cells, and inhibits acute heat stress-induced skin cell apoptosis via activation of AKT pathway. Our findings suggest that administration of allogeneic hucMSCs derived exosomes is an alternative approach for stem cell-based therapy and may represent a novel therapeutic strategy for skin injury repair.

CONCLUSION

Our results have clearly demonstrated that hucMSC-Ex enhances skin second-degree burn injury repair and

Wnt4 is the key mediator delivered by hucMSC-Ex in cutaneous wound healing.

ACKNOWLEDGMENTS

This work was supported by the Major Research Plan of the National Natural Science Foundation of China (Grant No. 91129718), the National Natural Science Foundation of China (Grant no. 31340040, 81272481, 81270214), Jiangsu Province for Outstanding Sci-tech Innovation Team in Colleges and Universities (Grant no. SJK2013-10), Jiangsu Province's Outstanding Medical Academic Leader and Sci-tech Innovation Team Program (Grant no. LJ201117), Jiangsu Province Doctoral Innovation Fund (Grant no. CXZZ13_0703).

DISCLOSURE OF POTENTIAL CONFLICTS OF INTEREST

None

AUTHOR CONTRIBUTIONS

B.Z.: Conception and design, Collection and/or assembly of data, Data analysis and interpretation, Manuscript writing; M.W.: Conception and design, Collection and/or assembly of data, Data analysis and interpretation, Manuscript writing; A.G.: Conception and design, Collection and/or assembly of data, Data analysis and interpretation, Manuscript writing; X.Z.: Collection and/or assembly of data; X.W.: Collection and/or assembly of data; Y.Z.: Collection and/or assembly of data; H.S.: Collection and/or assembly of data; L.W.: Conception and design, Data analysis; W.Z.: Conception and design, Data analysis; H.Q. and W.X.: Conception and design, Data analysis and interpretation, Financial support, Manuscript writing, Final approval of manuscript.

REFERENCES

- Martin P. Wound healing—aiming for perfect skin regeneration. *Science* 1997;276:76-81.
- Singer AJ, Clark RA. Cutaneous wound healing. *N Engl J Med* 1999;1:738-746.
- Bielefeld KA, Amini-Nik S, Alman BA. Cutaneous wound healing: recruiting developmental pathways for regeneration. *Cell Mol Life Sci* 2013;70: 2069-2081.
- Iqbal T, Saaiq M, Ali Z. Epidemiology and outcome of burns: early experience at the country's first national burns centre. *Burns* 2013;39:358-362.
- Blais M, Parenteau-Bareil R, Cadau S, et al. Concise review: tissue-engineered skin and nerve regeneration in burn treatment. *Stem Cells Transl Med* 2013;2:646-661.
- Phinney DG, Prockop DJ. Concise review: mesenchymal stem/multipotent stromal cells: the state of transdifferentiation and modes of tissue repair—current views. *Stem Cells* 2007;26:2896-2902.
- Kolf CM, Cho E, Tuan RS. Mesenchymal stromal cells. Biology of adult mesenchymal stem cells: regulation of niche, self-renewal and differentiation. *Arthritis Res Ther* 2007;9: 204.
- Panepucci RA, Siufi JL, Silva Wa, et al. Comparison of gene expression of umbilical cord vein and bone marrow-derived mesenchymal stem cells. *Stem Cells* 2004;22:1263-1278.
- Chen W, Liu J, Manuchehrabadi N, et al. Umbilical cord and bone marrow mesenchymal stem cell seeding on macroporous calcium phosphate for bone regeneration in rat cranial defects. *Biomaterials* 2013;34:9917-9925.
- Cao H, Qian H, Xu W, et al. Mesenchymal stem cells derived from human umbilical cord ameliorate ischemia/reperfusion-induced acute renal failure in rats. *Biotechnol Lett* 2010;32:726-732.
- Chen Y, Qian H, Zhu W, et al. Hepatocyte growth factor modification promotes the amelioration effects of human umbilical cord mesenchymal stem cells on rat acute kidney injury. *Stem Cells Dev* 2011;20:103-113.
- Qian H, Yang H, Xu W, et al. Bone marrow mesenchymal stem cells ameliorate rat acute renal failure by differentiation into renal tubular epithelial-like cells. *Int J Mol Med* 2008;22:326-332.
- Yan Y, Xu W, Qian H, et al. Mesenchymal stem cells from human umbilical cords ameliorate mouse hepatic injury in vivo. *Liver Int* 2009;29:366-366.
- Bruno S, Grange C, Deregibus MC, et al. Mesenchymal stem cell-derived microvesicles protect against acute tubular injury. *J Am Soc Nephrol* 2009;20:1063-1067.
- Gatti S, Bruno S, Deregibus MC, et al. Microvesicles derived from human adult mesenchymal stem cells protect against ischaemia-reperfusion-induced acute and chronic kidney injury. *Nephrol Dial Transplant* 2011;26:1474-1483.
- Xin H, Li Y, Buller B, et al. Exosome-mediated transfer of miR-133b from multipotent mesenchymal stromal cells to

neural cells contributes to neurite outgrowth. *Stem Cells* 2012;30:1666-1664.

17 Schorey JS, Bhatnagar S. Exosome function: from tumor immunology to pathogen biology. *Traffic* 2008;9:871-881.

18 Zhou Y, Xu H, Xu W, et al. Exosomes released by human umbilical cord mesenchymal stem cells protect against cisplatin-induced renal oxidative stress and apoptosis in vivo and in vitro. *Stem Cell Res Ther* 2013; 4:34.

19 Li T, Yan Y, Wang B, et al. Exosomes derived from human umbilical cord mesenchymal stem cells alleviate liver fibrosis. *Stem Cells Dev* 2013;22:846-864.

20 Huelsken J, Vogel R, Erdmann B, et al. Beta-catenin controls hair follicle morphogenesis and stem cell differentiation in the skin. *Cell* 2001;106:633-646.

21 Andl T, Reddy ST, Gaddapara T, et al. WNT signals are required for the initiation of hair follicle development. *Dev Cell* 2002;2:643-663.

22 Lim X, Tan SH, Koh WL, et al. Interfollicular epidermal stem cells self-renew via autocrine Wnt signaling. *Science* 2013;2:1226-1230.

23 Cheon SS, Wei Q, Gurung A, et al. Beta-catenin regulates wound size and mediates the effect of TGF-beta in cutaneous healing. *FASEB J* 2006;20:692-701.

24 Gross JC, Chaudhary V, Bartscherer K, et al. Active Wnt proteins are secreted on exosomes. *Nat Cell Biol* 2012;14:1036-1046.

25 Luga V, Zhang L, Vitoria-Petit AM, et al. Exosomes mediate stromal mobilization of autocrine Wnt-PCP signaling in breast cancer cell migration. *Cell* 2012;161:1642-1666.

26 Qiao C, Xu W, Zhu W, et al. Human mesenchymal stem cells isolated from the umbilical cord. *Cell Biol Int* 2008;32:8-16.

27 Nakamura Y, Ishikawa H, Kawai K, et al. Enhanced wound healing by topical administration of mesenchymal stem cells transfected with stromal cell-derived factor-1. *Biomaterials* 2013;34:9393-9400.

28 Morimoto N, Takemoto S, Kanda N, et al. The utilization of animal product-free media and autologous serum in an autologous dermal substitute culture. *J Surg Res* 2011;171:339-346.

29 Nacer Khodja A, Mahlous M, Tahtat D, et al. Evaluation of healing activity of PVA/chitosan hydrogels on deep second degree burn: pharmacological and toxicological tests. *Burns* 2013;39:98-104.

30 Wu Y, Chen L, Scott PG, et al. Mesenchymal stem cells enhance wound healing through differentiation and angiogenesis. *Stem Cells* 2007;26:2648-2669.

31 Chen YG, Li Z, Wang XF. Where PI3K/AKT meets Smads: the crosstalk determines human embryonic stem cell fate. *Cell Stem Cell* 2012;10:231-232.

32 Chen EY, Mazure NM, Cooper JA, et al. Hypoxia activates a platelet-derived growth factor receptor/phosphatidylinositol 3-kinase/Akt pathway that results in glycogen synthase kinase-3 inactivation. *Cancer Res* 2001;61:2429-2433.

33 Valadi H, Ekström K, Bossios A, et al. Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nat Cell Biol* 2007;9:664-669.

34 Vella LJ, Sharples RA, Nisbet RM, et al. The role of exosomes in the processing of proteins associated with neurodegenerative diseases. *Eur Biophys J* 2008;37:323-332.

35 Herrera MB, Fonsato V, Gatti S, et al. Human liver stem cell-derived microvesicles accelerate hepatic regeneration in hepatectomized rats. *J Cell Mol Med* 2010;14:1606-1618.

36 Peinado H, Alečković M, Lavotshkin S, et al. Melanoma exosomes educate bone marrow progenitor cells toward a pro-metastatic phenotype through MET. *Nat Med* 2012;18:883-891.

37 Atay S, Banskota S, Crow J, et al. Oncogenic KIT-containing exosomes increase gastrointestinal stromal tumor cell invasion. *PNAS* 2014;111:711-716.

38 Takada R1, Satomi Y, Kurata T, et al. Monounsaturated Fatty Acid Modification of Wnt Protein: Its Role in Wnt Secretion. *Developmental Cell* 2006;11: 791-801.

39 Doubravska L, Krausova M, Gradl D, et al. Fatty acid modification of Wnt1 and Wnt3a at serine is prerequisite for lipidation at cysteine and is essential for Wnt signalling. *Cellular Signalling* 2011;23:837-848.

40 Coudreuse D, Korswagen HC. The making of Wnt: new insights into Wnt maturation, sorting and secretion. *Development* 2007;1:3-12.

41 Willert K, Nusse R. Wnt proteins. *Cold Spring Harb Perspect Biol* 2012;4:a007864.

42 Zhai L, Chaturvedi D, Cumberledge S. *Drosophila wnt-1* undergoes a hydrophobic modification and is targeted to lipid rafts, a process that requires porcupine. *J Biol Chem* 2004;279:33220-33207.

43 Willert K, Brown JD, Danenberg E, et al. Wnt proteins are lipid-modified and can act as stem cell growth factors. *Nature* 2003;423:448-52.

44 Schulte G, Bryja V, Rawal N, et al. Purified Wnt-5a increases differentiation of mid-brain dopaminergic cells and dishevelled phosphorylation. *J Neurochem* 2005;92:1550-1553.

45 Mikels AJ, Nusse R. Wnts as ligands: processing, secretion and reception. *Oncogene* 2006;25:7461-7468.

46 Korkut C, Ataman B, Ramachandran P, et al. Trans-synaptic transmission of vesicular Wnt signals through Evi/Wntless. *Cell* 2009;139:393-404.

47 Taelman VF, Dobrowolski R, Plouhinec JL, et al. Wnt signaling requires sequestration of glycogen synthase kinase 3 inside multivesicular endosomes. *Cell* 2010;143:1136-1148.

48 Dittmer J, Leyh B. Paracrine effects of stem cells in wound healing and cancer progression (Review). *Int J Oncol* 2014;44:1789-1798.

49 Fuchs Y, Brown S, Gorenc T, et al. Sept4/ARTS regulates stem cell apoptosis and skin regeneration. *Science* 2013;341:286-289.

50 Banerjee Mustafi S, Chakraborty PK, Dey RS, et al. Heat stress upregulates chaperone heat shock protein 70 and antioxidant manganese superoxide dismutase through reactive oxygen species (ROS), p38MAPK, and AKT. *Cell Stress Chaperones* 2009;14:679-689.

51 Deregibus MC, Cantaluppi V, Calogero R, et al. Endothelial progenitor cell derived microvesicles activate an angiogenic program in endothelial cells by a horizontal transfer of mRNA. *Blood* 2007;110:2440-2448.



See www.StemCells.com for supporting information available online. STEM CELLS ; 00:000-000

Figure 1. HucMSC-Ex accelerated the recovery of skin second-deep burn injury in rats. (A) Representative micrographs of wound histological images (H&E stain) at 1 week and 2 weeks after treatment. (B) Representative images of immunohistochemical staining of PCNA in each group. (C) Wound histological scores were calculated at 1 week and 2 weeks after treatment (n=6; * p<0.05, *** p<0.001). (D) Quantitative analyses for relative mRNA level of type I and III collagen in wound tissue at 1 week and 2 weeks after treatment (n=6; **p<0.01, *** p<0.001). (E) Representative immunofluorescence images of CK19 expression showed re-epithelialization in wound area. Scale bar=100 μ m. (F) Western blot assay for PCNA and CK19 expression in wounds at 1 week and 2 weeks after treatment.




Figure 2. HucMSC-Ex inhibited heat stress-induced apoptosis of keratinocytes and promoted their proliferation. (A) HaCAT and DFL cells were cultured under treatment of PBS, HFL1-Ex or hucMSC-Ex for 24 h after 43°C 40 min heat stress and subjected to flow cytometric assay for apoptotic cells. (n=3; **p<0.01, ***p<0.001). (B) HaCAT and (C) DFL cells were cultured under treatment of PBS, HFL1-Ex or hucMSC-Ex for 24 h after 43°C 40 min heat stress. The number of cells at indicated time points was shown (n=3; * p<0.05, **p<0.01, ***p<0.001). (D) Western blot assay for PCNA, Bcl-2 and Bax expression in HaCAT cells treated with PBS, HFL1-Ex or hucMSC-Ex for 24 h after 43°C 40 min heat stress. (E) Western blot assay for PCNA, Bcl-2 and Bax expression in DFL cells treated with PBS, HFL1-Ex or hucMSC-Ex for 24 h after 43°C 40 min heat stress.




Figure 3. HucMSC-Ex activated β -catenin signaling to prompt wound healing. (A) 293T cells transfected with the TOP-Flash or FOP-Flash luciferase reporter were treated with PBS, HFL1-Ex, or hucMSC-Ex. The ratio between TOP-Flash and FOP-Flash luciferase activity were determined at 24 h after treatment ($n=3$, $***p<0.001$). (B) HaCAT cells were treated with PBS, HFL1-Ex, or hucMSC-Ex for 24 h. The nuclear translocation of β -catenin was determined by immunofluorescence staining. (C) Western blot assay for the expression of β -catenin downstream targets (cyclin D1, cyclin D3 and N-cadherin) in HaCAT cells treated with PBS or hucMSC-Ex in the presence or absence of ICG001(20 μ M/ml). (D) The ratio between Top-Flash and Fop-Flash luciferase activity was determined at 24 h after treatment with hucMSC-Ex in the presence or absence of ICG001 (20 μ M/ml) ($n=3$, $**P<0.01$, $***p<0.001$). (E) The migration of HaCAT cells treated with hucMSC-Ex for 12 and 24 h in the presence or absence of ICG001 (20 μ M/ml) was determined by cell scratch assay ($n=3$, $**P<0.01$, $***p<0.001$). (F) Cell cycle analysis of HaCAT cells treated with hucMSC-Ex for 24 h in the presence or absence of ICG001 (20 μ M/ml) ($n=3$, $***p<0.001$). (G) The rat model was subcutaneously injected with hucMSC-Ex and ICG001 (1 mg/rat). The wound was subjected to H&E staining and immunohistochemical staining for PCNA expression at 1 week after treatment. Scale bar=100 μ m. (H) Wound histological scores ($n=6$ at 1 week after treatment; $*** p<0.001$).




Figure 4. Wnt4 in hucMSC-Ex mediated the activation of β -catenin signaling and wound healing. (A) The mRNA levels of Wnt family members in hucMSC and HFL1 cell were detected by using quantitative RT-PCR ($n=3$, $*p<0.05$, $***p<0.001$). (B) Western blot assay for the expression of Wnt4 in hucMSC-Ex and HFL1-Ex. (C) HucMSCs were transfected with Wnt4-shRNA or GFP-shRNA by lentivirus. The expression of Wnt4 in hucMSC and hucMSC-Ex was determined by using Western blot assay. (D) The ratio between TOP-Flash and FOP-Flash luciferase activity was determined at 24 h after treatment with exosomes from hucMSC transfected with Wnt4-shRNA (Wnt4-shRNA-Ex) or GFP-shRNA (GFP-shRNA-Ex) by lentivirus. ($n=3$, $***p<0.001$). (E) HaCAT cells were treated with PBS, GFP-shRNA-Ex or Wnt4-shRNA-Ex under normal or heat stress conditions for 24 h. The nuclear translocation of β -catenin was detected by immunofluorescence. (F) Western blot assay for the expression of β -catenin and its downstream targets (cyclin D1, cyclin D3 and N-cadherin) and the phosphorylation of GSK3 β in HaCAT cells treated with PBS, GFP-shRNA-Ex or Wnt4-shRNA-Ex under heat stress condition. (G) The migration of HaCAT cells treated with PBS, GFP-shRNA-Ex or Wnt4-shRNA-Ex for 12 and 24 h was detected by using cell scratch assay ($n=3$, $***p<0.001$). (H) Cell cycle transition analysis of HaCAT cells treated with PBS, GFP-shRNA-Ex or Wnt4-shRNA-Ex for 24 h ($n=3$, $*p<0.05$, $***p<0.001$). (I) Rat wound models were treated with PBS, GFP-shRNA-Ex or Wnt4-shRNA-Ex and then subjected to H&E staining and immunohistochemical staining of PCNA at 1 week after treatment. Scale bar=100 μ m. (J) Wound histological scores ($n=6$ at 1 week after treatment; $*** p<0.001$).




Figure 5. HucMSC-Ex inhibited heat stress-induced apoptosis by activating AKT signaling. (A) HaCAT and DFL cells were subjected to heat stress (43 °C, 40 min) and treated with HFL1-Ex or hucMSC-Ex for 12 h. The expression of total and phosphorylated AKT was determined by Western blot. (B) HaCAT cells were treated as described in (A) in the presence or absence of LY294002. The expression of total and phosphorylated AKT (p-AKT), Bcl-2 and Bax was determined by Western blot. (C) Western blot results examined the expression of p-AKT, Bcl-2 and Bax *in vivo*. (D) Representative immunofluorescence images of p-AKT expression in the injured skin tissues treated with hucMSC-Ex in the presence or absence of LY294002 (0.5mg/rat), HFL1-Ex and PBS. (E) TUNEL assay of apoptotic cells in the injured skin tissues treated with hucMSC-Ex in the presence or absence of LY294002 (0.5mg/rat), HFL1-Ex and PBS. (n=6, ***p<0.001). Scale bar=50 μ m. (F) Cell cycle distribution analysis of HaCAT cells treated with hucMSC-Ex in the presence or absence of LY294002 for 24 h (n=3, *p<0.05, **P<0.01, ***p<0.001). (G) The migratory ability of HaCAT cells treated with hucMSC-Ex in the presence or absence of LY294002 (50 μ M/ml) for 12 h and 24 h was examined by cell scratch assay (n=3, *p<0.05, **P<0.01, ***p<0.001).




Figure 6. HucMSC-Ex mediated bifurcated activation of AKT and Wnt4/ β -catenin signaling to promote wound healing. (A) The ratio between TOP-Flash and FOP-Flash luciferase activity was determined at 24 h after treatment with hucMSC-Ex in the presence or absence of LY294002 (50 μ M/ml) (n=3, **p<0.01). (B) HaCAT and DFL cells were treated with hucMSC-Ex in the presence or absence of LY294002 (50 μ M/ml) for 24 h. The nuclear translocation of β -catenin was detected by immunofluorescence. (C) Western blot assay for β -catenin and p-AKT and total AKT, p-GSK3 β and GSK3 β in DFL cells treated with PBS or hucMSC-Ex in the presence or absence of 50 μ M/ml of LY294002. (D) Immunofluorescence staining of p-AKT and β -catenin in consecutive sections of representative wounds that were treated with PBS, HFL1-Ex or hucMSC-Ex in the presence or absence of LY294002 (0.5 mg/rat). Scale bar=100 μ m. (E) Western blot assay for p-AKT, Bcl-2, and Bax expression in DFL cells treated with PBS or hucMSC-Ex in the presence or absence of ICG001. (F) Western blot assay for p-AKT, Bcl-2, and Bax in DFL cells treated with PBS, GFP-shRNA-Ex or Wnt4-shRNA-Ex under heat stress condition.

